

**CYCLOPENTOLATE HYDROCHLORIDE- cyclopentolate hydrochloride solution/ drops
Akorn, Inc.**

Cyclopentolate Hydrochloride Ophthalmic Solution, USP -Sterile

CC00N Rev. 10/16

Rx only

DESCRIPTION:


Cyclopentolate Hydrochloride Ophthalmic Solution is an anticholinergic prepared as a sterile, borate buffered solution for topical ocular use. It is supplied in two strengths.

Chemical name:

2-(Dimethylamino)ethyl 1-hydroxy- α -phenylcyclopentaneacetate hydrochloride

MW=327.85 C₁₇H₂₅NO₃ \cdot HCl

The active ingredient is represented by the structural formula:



Cyclopentolate Hydrochloride Ophthalmic Solution USP, 1% & 2%

Each mL contains: Active: Cyclopentolate Hydrochloride 10 mg (1%) or 20 mg (2%).

Inactives: Boric Acid, Edetate Disodium, Potassium Chloride (except 2% strength), Sodium Carbonate and/or Hydrochloric Acid may be added to adjust pH (3.0 to 5.5) and Water for Injection.

Preservative: Benzalkonium Chloride 0.1 mg (0.01%).

CLINICAL PHARMACOLOGY:

This anticholinergic preparation blocks the responses of the sphincter muscle of the iris and the accommodative muscle of the ciliary body to cholinergic stimulation, producing pupillary dilation (mydriasis) and paralysis of accommodation (cycloplegia). It acts rapidly, but has a shorter duration than atropine. Maximal cycloplegia occurs within 25 to 75 minutes after instillation. Complete recovery of accommodation usually takes 6 to 24 hours. Complete recovery from mydriasis in some individuals may require several days. Heavily pigmented irides may require more doses than lightly pigmented irides.

INDICATIONS AND USAGE:

Cyclopentolate hydrochloride ophthalmic solution is used to produce mydriasis and cycloplegia.

CONTRAINDICATIONS:

Cyclopentolate Hydrochloride Ophthalmic Solution should not be used when untreated narrow-angle glaucoma, or untreated anatomically narrow angles are present, or if the patient is hypersensitive to any component of this preparation.

WARNINGS:

For topical ophthalmic use only. Not for injection. This preparation may cause CNS disturbances. This is especially true in younger age groups, but may occur at any age, especially with the stronger solutions. Infants are especially prone to CNS and cardiopulmonary side effects from cyclopentolate. To minimize absorption, use only 1 drop of 0.5% cyclopentolate hydrochloride ophthalmic solution per eye, followed by pressure applied over the nasolacrimal sac for two to three minutes. Observe infants closely for at least 30 minutes.

Mydriatics may produce a transient elevation of intraocular pressure.

PRECAUTIONS:

General: The lacrimal sac should be compressed by digital pressure for two to three minutes after instillation to reduce excessive systemic absorption. Caution should be observed when considering use of this medication in the presence of Down's syndrome and in those predisposed to angle-closure glaucoma.

Information for Patients: Do not touch dropper tip to any surface, as this may contaminate the solution. A transient burning sensation may occur upon instillation. Patients should be advised not to drive or engage in other hazardous activities while pupils are dilated. Patients may experience sensitivity to light and should protect eyes in bright illumination during dilation. Parents should be warned not to get this preparation in their child's mouth and to wash their own hands and the child's hands following administration. Feeding intolerance may follow ophthalmic use of this product in infants. It is recommended that feeding be withheld for four (4) hours after examination.

Drug Interactions: Cyclopentolate may interfere with the ocular anti-hypertensive action of carbachol, pilocarpine, or ophthalmic cholinesterase inhibitors.

Carcinogenesis, Mutagenesis, and Impairment of Fertility: Studies in animals or humans have not been conducted to evaluate the carcinogenic potential of cyclopentolate.

Pregnancy: Pregnancy Category C. Animal reproduction studies have not been conducted with cyclopentolate. It is also not known whether cyclopentolate can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Cyclopentolate should be administered to a pregnant woman only if clearly needed.

Nursing Mothers: It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when cyclopentolate hydrochloride is administered to a nursing woman.

Pediatric Use: Use of cyclopentolate has been associated with psychotic reactions and behavioral disturbances in pediatric patients. Increased susceptibility to cyclopentolate has been reported in infants, young children, and in children with spastic paralysis or brain damage. These disturbances include ataxia, incoherent speech, restlessness, hallucinations, hyperactivity, seizures, disorientation as to time and place, and failure to recognize people. Feeding intolerance may follow ophthalmic use of this product in infants. It is recommended that feeding be withheld for four (4) hours after examination. Observe infants closely for at least 30 minutes (see **WARNINGS**).

ADVERSE REACTIONS:

Ocular: Increased intraocular pressure, burning, photophobia, blurred vision, irritation, hyperemia, conjunctivitis, blepharoconjunctivitis, punctate keratitis, synechiae have been reported. **Non-ocular:** Use of cyclopentolate has been associated with psychotic reactions and behavioral disturbances, usually in children, especially with 2% concentration. These disturbances include ataxia, incoherent speech, restlessness, hallucinations, hyperactivity, seizures, disorientation as to time and place, and failure to recognize people. This drug produces reactions similar to those of other anticholinergic drugs, but the central nervous system manifestations as noted above are more common. Other

manifestations of anticholinergic drugs are skin rash, abdominal distention in infants, unusual drowsiness, tachycardia, hyperpyrexia, vasodilation, urinary retention, diminished gastrointestinal motility and decreased secretion in salivary and sweat glands, pharynx, bronchi and nasal passages. Severe manifestations of toxicity include coma, medullary paralysis and death.

OVERDOSAGE:

Excessive dosage may produce behavioral disturbances, tachycardia, hyperpyrexia, hypertension, elevated intraocular pressure, vasodilation, urinary retention, diminished gastrointestinal motility and decreased secretion in salivary and sweat glands, pharynx, bronchi and nasal passages. Patients exhibiting signs of overdose should receive supportive care and monitoring.

DOSAGE AND ADMINISTRATION:

Adults: Instill one or two drops of 1% or 2% solution in the eye which may be repeated in five to ten minutes if necessary. Complete recovery usually occurs in 24 hours. Complete recovery from mydriasis in some individuals may require several days. **Children:** Instill one or two drops of 1% or 2% solution in the eye which may be repeated five to ten minutes later by a second application of 1% solution if necessary.

HOW SUPPLIED:

Cyclopentolate Hydrochloride Ophthalmic Solution, USP is a sterile ophthalmic solution supplied in white opaque plastic dropper bottles as follows:

Cyclopentolate Hydrochloride Ophthalmic Solution USP, 1%
2 mL NDC 17478-100-02
5 mL NDC 17478-100-10
15 mL NDC 17478-100-12

Cyclopentolate Hydrochloride Ophthalmic Solution USP, 2%
2 mL NDC 17478-097-02
5 mL NDC 17478-097-10
15 mL NDC 17478-097-12

DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING.

Storage: Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Keep tightly closed.

KEEP OUT OF THE REACH OF CHILDREN.

AKORN
Manufactured by: Akorn, Inc.
Lake Forest, IL 60045
CC00N
Rev. 10/16

Principal Display Panel Text for Container Label:

NDC 17478-100-02
Cyclopentolate
Hydrochloride
Ophthalmic
Solution, USP

1%

For Topical

Ophthalmic Use Only

Rx only 2 mL

EXP. LOT

(01)00317478100027

Storage: Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Keep tightly closed.

KEEP OUT OF REACH OF CHILDREN.

NDC 17478-100-02

Cyclopentolate Hydrochloride Ophthalmic Solution, USP

1%

For Topical Ophthalmic Use Only

Rx only 2 mL

Each mL contains:
Active: Cyclopentolate Hydrochloride 10 mg (1%).

PRECAUTION: Do not touch dropper tip to any surface, as this may contaminate the solution.

Mfd. by: **Akorn, Inc.**
Lake Forest, IL 60045

CCADL Rev. 05/17

Principal Display Panel Text for Carton Label:

NDC 17478-100-02

Cyclopentolate

Hydrochloride

Ophthalmic

Solution, USP

1%

For Topical

Ophthalmic Use Only

2 mL

Rx only Akorn logo



Principal Display Panel Text for Container Label:

NDC 17478-097-10

Cyclopentolate

Hydrochloride

Ophthalmic

Solution, USP

2%

5 mL

Rx only Sterile

EXP. LOT
(01)00317478097105

Each mL contains:
Active: Cyclopentolate Hydrochloride 20 mg (2%);
Inactives: Boric Acid, Edetate Disodium, Sodium Carbonate and/or Hydrochloric Acid may be added to adjust pH (3.0 to 5.5) and Water for Injection;
Preservative: Benzalkonium Chloride 0.1 mg (0.01%).
Mfd. by: **Akorn, Inc.**
Lake Forest, IL 60045

NDC 17478-097-10
Cyclopentolate Hydrochloride Ophthalmic Solution, USP
2 %
5 mL
Rx only Sterile

For Topical Ophthalmic Use Only.
Usual Dosage: See package insert for dosage information.
Storage: Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Keep tightly closed.
PRECAUTION: Do not touch dropper tip to any surface, as this may contaminate the solution.
KEEP OUT OF REACH OF CHILDREN.
CCBEL Rev. 10/16

Principal Display Panel Text for Carton Label:

NDC 17478-097-10

Cyclopentolate

Hydrochloride

Ophthalmic

Solution, USP

2%

5 mL

Sterile

Rx only Akorn logo



CCBEC0417

NDC 17478-097-10

NDC 17478-097-10

Each mL contains:

Active: Cyclopentolate Hydrochloride 20 mg (2%);
Inactives: Boric Acid, Edetate Disodium, Sodium Carbonate and/or Hydrochloric Acid may be added to adjust pH (3.0 to 5.5) and Water for Injection;
Preservative: Benzalkonium Chloride 0.1 mg (0.01%).
Note: Bottle filled to 1/2 capacity for proper drop control.

Cyclopentolate Hydrochloride Ophthalmic Solution, USP

2%

For Topical Ophthalmic Use Only.

Usual Dosage: One drop in each eye followed by a second drop in five minutes if necessary. See package insert for dosage information.

Storage: Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature]. Keep tightly closed.

Cyclopentolate Hydrochloride Ophthalmic Solution, USP

2%

PRECAUTION: Do not touch dropper tip to any surface, as this may contaminate the solution.

DO NOT USE IF IMPRINTED SEAL IS BROKEN OR MISSING.

KEEP OUT OF REACH OF CHILDREN.

5 mL

Sterile



Manufactured by:
Akorn, Inc.
Lake Forest, IL 60045

CCBEC Rev. 04/17

5 mL

Sterile

Rx only **AKORN**

Rx only **AKORN**

CYCLOPENTOLATE HYDROCHLORIDE

cyclopentolate hydrochloride solution/ drops

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:17478-100
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Cyclopentolate Hydrochloride (UNII: 736I6971TE) (Cyclopentolate - UNII:I76F4SHP7J)	Cyclopentolate Hydrochloride	10 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
benzalkonium chloride (UNII: F5UM2KM3W7)	
boric acid (UNII: R57ZHV85D4)	
edetate disodium (UNII: 7FLD91C86K)	
potassium chloride (UNII: 660YQ98I10)	
sodium carbonate (UNII: 45P3261C7T)	
hydrochloric acid (UNII: QTT17582CB)	
water (UNII: 059QF0KO0R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17478-100-02	1 in 1 CARTON	01/13/1997	
1		2 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		
2	NDC:17478-100-10	1 in 1 CARTON	01/13/1997	
2		5 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		
3	NDC:17478-100-12	1 in 1 CARTON	01/13/1997	
3		15 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040164	01/13/1997	

CYCLOPENTOLATE HYDROCHLORIDE

cyclopentolate hydrochloride solution/ drops

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:17478-097
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
Cyclopentolate Hydrochloride (UNII: 736I6971TE) (Cyclopentolate -	Cyclopentolate	20 mg in 1 mL

UNII:I76F4SHP7J)	Hydrochloride	20 mg in 1 mL
------------------	---------------	---------------

Inactive Ingredients

Ingredient Name	Strength
benzalkonium chloride (UNII: F5UM2KM3W7)	
boric acid (UNII: R57ZHV85D4)	
edetate disodium (UNII: 7FLD91C86K)	
sodium carbonate (UNII: 45P3261C7T)	
hydrochloric acid (UNII: QTT17582CB)	
water (UNII: 059QF0K00R)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:17478-097-02	1 in 1 CARTON	01/13/1997	
1		2 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		
2	NDC:17478-097-10	1 in 1 CARTON	01/13/1997	
2		5 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		
3	NDC:17478-097-12	1 in 1 CARTON	01/13/1997	
3		15 mL in 1 BOTTLE, DROPPER; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA040165	01/13/1997	

Labeler - Akorn, Inc. (062649876)

Establishment

Name	Address	ID/FEI	Business Operations
Akorn, Inc.		155135783	MANUFACTURE(17478-100, 17478-097) , REPACK(17478-100, 17478-097) , ANALYSIS(17478-100, 17478-097)

Revised: 8/2017

Akorn, Inc.