


ALCAINE - proparacaine hydrochloride solution/ drops
Alcon Laboratories, Inc.

Alcaine®
(proparacaine hydrochloride
ophthalmic solution, USP) 0.5%

DESCRIPTION:

ALCAINE® (proparacaine hydrochloride ophthalmic solution, USP) 0.5% is a topical local anesthetic for ophthalmic use. The active ingredient is represented by the structural formula:



Established Name: Proparacaine Hydrochloride

Chemical Name: Benzoic acid, 3-amino-4-propoxy-,2-(diethylamino) ethyl ester, monohydrochloride.

Molecular Weight: 330.85 g/mol

Each mL contains of ALCAINE® (proparacaine hydrochloride ophthalmic solution, USP) 0.5%:

Active: proparacaine hydrochloride 5 mg 0.5%. **Preservative:** benzalkonium chloride (0.01%).

Inactives: glycerin and purified water. The pH may be adjusted with hydrochloric acid and/or sodium hydroxide.

CLINICAL PHARMACOLOGY:

ALCAINE® ophthalmic solution is a rapidly-acting topical anesthetic, with induced anesthesia lasting approximately 10-20 minutes.

INDICATIONS AND USAGE:

ALCAINE® ophthalmic solution is indicated for procedures in which a topical ophthalmic anesthetic is indicated: corneal anesthesia of short duration, e.g. tonometry, gonioscopy, removal of corneal foreign bodies, and for short corneal and conjunctival procedures.

CONTRAINDICATIONS:

ALCAINE® ophthalmic solution should be considered contraindicated in patients with known hypersensitivity to any of the ingredients of this preparation.

WARNINGS:

NOT FOR INJECTION - FOR TOPICAL OPHTHALMIC USE ONLY. Prolonged use of a topical ocular anesthetic is not recommended. It may produce permanent corneal opacification with accompanying visual loss.

PRECAUTIONS:

Carcinogenesis, Mutagenesis, Impairment of Fertility.

Long-term studies in animals have not been performed to evaluate carcinogenic potential, mutagenicity, or possible impairment of fertility in males or females.

Pregnancy:

Animal reproduction studies have not been conducted with ALCaine® (proparacaine hydrochloride ophthalmic solution, USP) 0.5%. It is also not known whether proparacaine hydrochloride can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. Proparacaine hydrochloride should be administered to a pregnant woman only if clearly needed.

Nursing Mothers:

It is not known whether this drug is excreted in human milk. Because many drugs are excreted in human milk, caution should be exercised when proparacaine hydrochloride is administered to a nursing woman.

Pediatric Use:

Safety and effectiveness of proparacaine hydrochloride ophthalmic solution in pediatric patients have been established. Use of proparacaine hydrochloride is supported by evidence from adequate and well-controlled studies in adults and children over the age of twelve, and safety information in neonates and other pediatric patients.

Geriatric Use:

No overall clinical differences in safety or effectiveness have been observed between the elderly and other adult patients.

ADVERSE REACTIONS:

Occasional temporary stinging, burning and conjunctival redness may occur with the use of proparacaine. A rare, severe, immediate-type, apparently hyperallergic corneal reaction characterized by acute, intense and diffuse epithelial keratitis, a gray, ground glass appearance, sloughing of large areas of necrotic epithelium, corneal filaments and, sometimes, iritis with descemetitis has been reported.

Allergic contact dermatitis from proparacaine with drying and fissuring of the fingertips has also been reported.

DOSAGE AND ADMINISTRATION:

Usual Dosage: Removal of foreign bodies and sutures, and for tonometry: 1 to 2 drops (in single instillations) in each eye before operating.

Short corneal and conjunctival procedures: 1 drop in each eye every 5 to 10 minutes for 5 to 7 doses.

NOTE: ALCaine® (proparacaine hydrochloride ophthalmic solution, USP) 0.5% should be clear to straw-color. If the solution becomes darker, discard the solution.

HOW SUPPLIED:

ALCaine® (proparacaine hydrochloride ophthalmic solution, USP) 0.5% is supplied in 15 mL DROP-

TAINER® dispensers as follows:

15 mL NDC 0998-0016-15

Storage:

Bottle must be stored in unit carton to protect contents from light.

Store bottles under refrigeration at 2°C to 8°C (36°F-46°F).

Rx Only

Distributed by:

ALCON LABORATORIES, INC.

Fort Worth, Texas 76134

Revised: April 2018

9016949-1118

PRINCIPAL DISPLAY PANEL

NDC 0998-0016-15

Alcon

Alcaine®

(proparacaine hydrochloride ophthalmic solution USP) 0.5%

15 mL Sterile

Rx Only

PRECAUTION: NOT FOR INJECTION. FOR TOPICAL OPHTHALMIC USE ONLY. Do not touch dropper tip to any surface, as this may contaminate the solution.

USUAL DOSAGE: 1 or 2 drops. Read enclosed insert.

Bottles must be stored in unit carton to protect from light.

STORAGE: Store between 2°-8°C (36°-46°F).

ALCAINE® (proparacaine hydrochloride ophthalmic solution) should be clear to straw-color. If the solution becomes darker, discard the solution.

INGREDIENTS: Each mL contains: Active: proparacaine hydrochloride 5 mg (0.5%). **Preservative:** benzalkonium chloride 0.01%. **Inactives:** glycerin, hydrochloric acid and/or sodium hydroxide (to adjust pH), purified water

pH range 4.0 to 6.0

1-800-757-9195

MedInfo@AlconLabs.com

Alcon

ALCON LABORATORIES, INC.

6201 South Freeway

Fort Worth, Texas 76134 USA

A Novartis Division
Printed in USA
9016948- 0019

SN:

LOT:

EXP:

GTIN: 00309980016156



NDC 0998-0016-15

Alcon

Alcaine®
(proparacaine hydrochloride ophthalmic solution USP) 0.5%

Sterile 15 mL

Rx Only

PRECAUTION: NOT FOR INJECTION. FOR TOPICAL OPHTHALMIC USE ONLY. Do not touch dropper tip to any surface as this may contaminate the solution.

USUAL DOSAGE: 1 or 2 drops. Read enclosed insert.

STORAGE: Store between 2°-8°C (36°-46°F).

Bottles must be stored in unit carton to protect from light.


ALCAINE® (proparacaine hydrochloride ophthalmic solution) should be clear to straw-color. If the solution becomes darker, discard the solution.

INGREDIENTS: Each mL contains: Active: proparacaine hydrochloride 5 mg (0.5%).

ALCON LABORATORIES, INC.
Fort Worth, Texas 76134 USA Printed in USA
A Novartis Division

LOT/EXP.:

H15466-1118



ALCAINE

proparacaine hydrochloride solution/ drops

Product Information

Product Type	HUMAN PRESCRIPTION DRUG	Item Code (Source)	NDC:0998-0016
Route of Administration	OPHTHALMIC		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
PROPARACAINE HYDROCHLORIDE (UNII: U96OL57GOY) (PROPARACAINE - UNII:B4OB0JH1X)	PROPARACAINE HYDROCHLORIDE	5 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7)	
GLYCERIN (UNII: PDC6A3C0OX)	
WATER (UNII: 059QF0K00R)	
HYDROCHLORIC ACID (UNII: QTT17582CB)	
SODIUM HYDROXIDE (UNII: 55X04QC32I)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0998-0016-15	1 in 1 CARTON	10/19/1973	
1		15 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA080027	10/19/1973	

Labeler - Alcon Laboratories, Inc. (008018525)

Registrant - Alcon Laboratories, Inc. (008018525)

Establishment

Name	Address	ID/FEI	Business Operations
Alcon Research, LLC		007672236	manufacture(0998-0016)